

Documents et textes de références

- "Recommandations pour la prise en charge de MRSA dans les EMS genevois" - Planification sanitaire Qualitative - Genève
- "Mesures pour la prévention des maladies transmissibles à l'hôpital" - Unité d'épidémiologie hospitalière / Institut central des hôpitaux valaisans – Valais
- Les mesures de contrôle et prévention des infections à *Staphylococcus aureus* résistant à la méthicilline (SARM) au Québec, édité par la Direction des communications du ministère de la Santé et des Services sociaux, 2000. www.msss.gouv.qc.ca
- CDC, Part II. Recommendations for Isolation Precautions in Hospitals
- SHEA Guideline for preventing nosocomial transmission of multidrug-resistant strains of *Staphylococcus aureus* and Enterococcus. *Infect Control Hosp Epidemiol.* 2003; 24:362-386

Bibliographie

- MA Abramson and coll: Nosocomial methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* primary bacteria: at what costs? *Infect Control Hosp Epidemiol.* 1999; 20:408-11
- DP Calfee and coll. Spread of methicillin-resistant *Staphylococcus aureus* (MRSA) among household contacts of individuals with nosocomially acquired MRSA. *Infect Control Hosp Epidemiol.* 2003; 24: 422-426
- C Chaix and coll. Control of endemic methicillin-resistant *Staphylococcus aureus* : a cost-benefit analysis in an intensive care unit. *JAMA.* 1999;282:1745-1751
- BS Cooper and coll. Systematic review of isolation policies in the hospital management of methicillin-resistant *Staphylococcus aureus* : a review of the literature with epidemiological and economic modeling. *Health Technol Assess.* 2003;7:1-194
- BS Cooper and coll. Isolation measures in the hospital management of methicillin-resistant *Staphylococcus aureus* (MRSA): systematic review of the literature. *BMJ.* 2004;329-533
- BS Cooper and coll. Methicillin-resistant *Staphylococcus aureus* in hospitals and in the community : stealth dynamics and control catastrophes. *Proc Natl Acad Sci USA* 2004; 101:10223-10228
- SE Cosgrove and coll. Comparaison of mortality associated with methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* bacteremia. A meta-analysis. *Clin Infect Dis.* 2003; 36 :53-59
- SE Cosgrove and coll. The impact of antimicrobial resistance and economic outcomes. *Clin Infect Dis.* 2003 ;36 :1433-1437
- S Harbarth and coll. Randomized, placebo-controlled, double-blind trial to evaluate the efficacy of mupirocin for eradicating carriage of methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 1999; 43:1412-1416
- WR Jarvis. Controlling antimicrobial-resistant pathogens. *Infect Control Hosp Epidemiol.* 2004;25:369-372
- JA Jernigan and coll. Prevalence of and risk factors for colonization with methicillin-resistant *Staphylococcus aureus* in an outpatient clinic population. *Infect Control Hosp Epidemiol.* 2003;24:445-450
- TB Karchmer and coll. Cost-effectiveness of active surveillance cultures and contact/droplet precautions for control of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect.* 2002;51:126-132
- J Kluytmans. Reduction of surgical site infections in major surgery by elimination of nasal carriage of *Staphylococcus aureus*. *J Hosp Infect.* 1998;40 Suppl B:S25-29
- JC Lucet and coll. Prevalence and risk factors for carriage of methicillin-resistant *Staphylococcus aureus* at admission to the intensive care unit: results of a multicenter study. *Arch Intern Med.* 2003;163:181-188
- CG McHugh and coll. Risk factors and costs associated with methicillin-resistant bloodstream infections. *Infect Control Hosp Epidemiol.* 2004; 25:425-430
- G Papia and coll. Screening high-risk patients for methicillin-resistant strains of *Staphylococcus aureus* on admission to the hospital: is it cost effective? *Infect Control Hosp Epidemiol.* 1999;20:473-477
- CD Salgado and coll. Interventions to prevent methicillin-resistant *Staphylococcus aureus* transmission in health care facilities: what works? *Clin Microbiol Newsl.* 2003; 25:137-144
- V Tonic and coll. Comprehensive strategy to prevent nosocomial spread of methicillin-resistant *Staphylococcus aureus* in a highly endemic setting. *Arch Intern Med.* 2004; 164: 2038-2043
- MR Vriens and coll. Is methicillin-resistant *Staphylococcus aureus* more contagious than methicillin-susceptible *S. aureus* in a surgical intensive care unit? *Infect Control Hosp Epidemiol.* 2002;23:491-494
- HF Wertheim and coll. Low prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) at hospital admission in the Netherlands: the value of search and destroy and restrictive antibiotic use. *J Hosp Infect.* 2004;56:321-325